



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

1875

1875

THE  
HISTORY OF THE  
CITY OF BOSTON  
FROM 1630 TO 1875  
BY  
JOHN H. COVINGTON  
BOSTON  
1875

1875

1875

# ELEMENTARY MATHEMATICAL WORKS

BY THE

Right Rev. J. W. COLENSO, D.D. Lord Bishop of Natal.

**ARITHMETIC** designed for the use of **SCHOOLS**: to which is added a Chapter on **DECIMAL COINAGE**. New Edition [1866], thoroughly revised, with the addition of Notes and Examination-Papers ..... 12mo. 4s. 6d.

**KEY** to **COLENSO'S ARITHMETIC** for **SCHOOLS**, containing full **SOLUTIONS** of all the **EXERCISES**. By the Rev. J. HUNTER, M.A. .... 12mo. 6s.

**COLENSO'S ELEMENTARY ARITHMETIC**, for the use of **NATIONAL, ADULT, and COMMERCIAL SCHOOLS**; consisting of a Text-Book of Arithmetic, accompanied by a Series of easy and amusing Examples, arranged progressively, in order of difficulty..... 18mo. 1s. 9d.; or with **ANSWERS**, 2s. 3d.

**COLENSO'S ELEMENTARY ARITHMETIC** for **NATIONAL SCHOOLS** is divided into **FIVE PARTS**, which are sold separately as follows:—

- I. Text-Book, price 6d.
- II. Examples, **PART I. Simple Arithmetic**, price 4d.
- III. Examples, **PART II. Compound Arithmetic**, price 4d.
- IV. Examples, **PART III. Fractions, Decimals, Duodecimals, &c.** price 4d.
- V. **ANSWERS** to the Examples, with **SOLUTIONS** of the more difficult Questions, price 1s.

**COLENSO'S ARITHMETICAL TABLES**, on a Card ...price 1d.

**THE ELEMENTS** of **ALGEBRA** designed for the use of **SCHOOLS**, **PART I.** containing the Simpler Parts of the Science, suitable for general School purposes, and as required for the attainment of an ordinary B.A. degree ..... 12mo. 4s. 6d.—**KEY**, 5s.

**HUNTER'S EXAMINATION-QUESTIONS** on **COLENSO'S ALGEBRA, PART I.** with Graduated Exercises, Problems, &c..... 12mo. 2s. 6d.

**COLENSO'S ELEMENTS** of **ALGEBRA** designed for the use of **SCHOOLS, PART II.** adapted to suit the wants of more advanced Students; containing the higher parts of the subject, with an **APPENDIX** of more difficult Miscellaneous Examples and Equation Papers ..... 12mo. 6s.—**KEY**, 5s.

**COLENSO'S MISCELLANEOUS EXAMPLES and EQUATION PAPERS** from **PARTS I. and II.** of the *Algebra*; with the **ANSWERS** ..... 12mo. 2s. 6d.

**COLENSO'S ELEMENTS** of **ALGEBRA**, adapted for the use of **NATIONAL and ADULT SCHOOLS**; containing numerous easy Examples and Questions under every Rule, with a Selection of Miscellaneous Exercises for Practice ..... 18mo. 1s. 6d.—**KEY**, 2s. 6d.

**COLENSO'S ELEMENTS** of **ALGEBRA**, adapted for the use of **TEACHERS and STUDENTS in the UNIVERSITY** ..... 8vo. 12s. 6d.—**KEY**, 12mo. 7s. 6d.

**PLANE TRIGONOMETRY, PART I.** comprising the Measurement of Lines and Angles, the Numerical Values of the Trigonometrical Ratios; with the Use of Logarithms and the Exponential Theorem ..... 12mo. 3s. 6d.—**KEY**, 3s. 6d.

**COLENSO'S PLANE TRIGONOMETRY, PART II.** comprising the Summation of Series, the Trigonometrical Solution of Equations, and Miscellaneous Problems 12mo. 2s. 6d.—**KEY**, 6s.... The **KEYS** to the Two **PARTS** of the Trigonometry, together, price 8s. 6d.

**THE ELEMENTS** of **EUCLID** (as studied in the Universities); with a large Collection of **GEOMETRICAL PROBLEMS**..... 18mo. 4s. 6d.; or with a **KEY**, 6s. 6d.

**COLENSO'S GEOMETRICAL PROBLEMS and KEY.** 18mo. 3s. 6d.

**COLENSO'S GEOMETRICAL PROBLEMS** separately, for the use of Schools where other Editions of *Euclid* may be employed..... 18mo. 1s.

London: **LONGMANS, GREEN, and CO.** Paternoster Row.





**KEY**

**TO THE**

**‘CONIC SECTIONS.’**

LONDON  
PRINTED BY SPOTTISWOODE AND CO.  
NEW-STREET SQUARE

K E Y  
TO  
HUNTER'S INTRODUCTION  
TO THE  
CONIC SECTIONS.

BY THE  
REV. JOHN HUNTER, M.A.

FORMERLY VICE-PRINCIPAL  
OF THE TRAINING COLLEGE, BATTERSEA.



LONDON:  
LONGMANS, GREEN, AND CO.  
1866.

183. g. 17.





# KEY


TO

## ‘CONIC SECTIONS.’




### EXERCISES [A].


1.  $3x - 2y = 4$ . When  $y = 0$ , then  $x = 1\frac{1}{3}$ ; and when  $x = 0$ , then  $y = -2$ . Therefore, take  $OA = 1\frac{1}{3}$ ,  $OB = 2$ . Then AB is the line required.




2.  $x + 3 = 4y$ . When  $y = 0$ , then  $x = -3$ ; and when  $x = 0$ , then  $y = \frac{3}{4}$ . Therefore, take  $OA = 3$ ,  $OB = \frac{3}{4}$ . Then AB is the line required.




3.  $4x + 5y = 6$ . When  $y = 0$ , then  $x = 1\frac{1}{2}$ ; and when  $x = 0$ , then  $y = 1\frac{1}{5}$ . Therefore, take  $OA = 1\frac{1}{2}$ ,  $OB = 1\frac{1}{5}$ . Then AB is the line required.




4.  $\frac{1}{4}y - \frac{1}{3}x = 5$ ; or  $3y - 4x = 60$ .  
When  $y=0$ , then  $x=-15$ ; and  
when  $x=0$ , then  $y=20$ . There-  
fore, take  $OA=15$ ,  $OB=20$ . Then  
 $AB$  is the line required.




5.  $3x + 4y = -5$ . When  $y=0$ ,  
then  $x=-1\frac{2}{3}$ ; and when  $x=0$ ,  
then  $y=-1\frac{1}{4}$ . Therefore, take  
 $OA=1\frac{2}{3}$ ,  $OB=1\frac{1}{4}$ . Then  $AB$  is  
the line required.




6.  $x + y = 0$ ; or  $y = -x$ . Here  
the coefficient of  $x$  is  $-1$  which is  
the tangent of  $135^\circ$ . Therefore,  
make  $TOX=135^\circ$ , and  $OT$  is the  
line required. For any point  $P$   
being taken in  $OT$  will make  $AP$   
 $= OA$ , or  $y = -x$ .




7.  $x\sqrt{3} - y = 8$ ; or  $y = \sqrt{3}x - 8$ .  
Here the coefficient of  $x$  is  $\sqrt{3}$ ,  
which is the tangent of  $60^\circ$ . There-  
fore, take  $OB=8$ , and on it describe  
the equilateral triangle  $OCB$ . The  
straight line  $BAT$  passing through  
the middle point of  $OC$  is the line  
required. For  $TAX = OAB = 90^\circ$   
 $- OBA = 60^\circ$ .




8.  $x = -2$ . This signifies that  
the abscissa of every point on the  
required line is  $= -2$ . There-  
fore, take  $OA=2$ , and the required  
line is  $AT$ , parallel to the axis  
of  $y$ .



9.  $\frac{4}{3}x + \frac{2}{3} = \frac{1}{2}y$ ; or  $y = \frac{8}{3}x + \frac{4}{3}$ . Here the tangent of the angle which the line makes with the axis of  $x$  is  $\frac{8}{3}$ ; therefore, take  $\frac{PM}{OM} = \frac{8}{3}$ , and join  $OP$ ; then take  $OB = 1\frac{1}{3}$ , and through  $B$  draw parallel to  $OP$  the required line  $AT$ .



10.  $6y + x\sqrt{5} + \sqrt{10} = 0$ ; or  $y = -\frac{\sqrt{5}}{6}x - \frac{\sqrt{10}}{6}$ .



Here the tangent being  $-\frac{\sqrt{5}}{6}$ , take  $OM=6$ ,  $PM=\sqrt{5}$ , and join  $OP$ ; then take  $OB=\frac{1}{3}\sqrt{10}$ , and through  $B$  draw parallel to  $OP$  the required line  $AT$ .

To find lines corresponding to  $\sqrt{5}$  and  $\sqrt{10}$ , we may take  $OE=1$ ,  $OD=2$ , and  $ED$  will be  $=\sqrt{5}$ ; and if  $OF$  be taken  $=3$ , the line  $EF$  will be  $=\sqrt{10}$ .


### EXERCISES [B].

1. Take  $ON, QN, =3, 5$ ;  $OM, PM, =5, 2$ ; the line  $PQ$  is that required.

From art. 15 we have

$$y+5 = \frac{2+5}{-5-3}(x-3); \text{ whence}$$


$$7x+8y=19.$$



2. Take  $OB=12$ ;  $OM, PM, =11$ ,  
7;  $PB$  is the required line. Its  
equation (by art. 15) is

$$y+12=\frac{7+12}{11-0}(x-0);$$

$$\text{or } 19x-11y=132.$$



3. The given line is  $y=\frac{3}{7}x-\frac{36}{7}$ . And, by art. 16, the  
equation to a line passing through the point  $(0, 0)$  parallel  
to the given line is  $y-0=\frac{3}{7}(x-0)$ , or  $3x=7y$ .


Also the equation to a line passing through the point  
 $(13, 4)$  parallel to the given line is

$$y-4=\frac{3}{7}(x-13); \text{ or } 3x-7y=11.$$

4. The given line is  
 $y=-\frac{5}{8}x+\frac{4}{3}$ . Take therefore

$OB=\frac{4}{3}$ , and  $\frac{PM}{OM}=\frac{5}{3}$ ; join  $OP$ ,

and through  $B$ , parallel to  $OP$ ,  
draw  $AB$ , which represents  
the given line.



Then take  $AN=5$ ,  $QN=1$ ,  
and the perpendicular  $QR$  on  
 $AB$  is the required line.

Now, by art. 18,  $y+1=\frac{3}{8}(x-5)$ ; or  $3x-5y=20$ .

5. The given line is  $y=-\frac{7}{18}x+14$ . And by art. 19 we  
have

$$y-30=\frac{-\frac{7}{18}+\sqrt{3}}{1+\frac{1}{18}\sqrt{3}}(x-5);$$

or, rationalising the denominators,

$$y-30=\frac{305\sqrt{3}+448}{+109}(x-5);$$

which gives for the required equations

$$109y+(448\pm 305\sqrt{3})x-5(1102\pm 305\sqrt{3})=0.$$

6. By art. 20,  $PQ^2 = (5+7)^2 + (-4-12)^2 = 12^2 + 16^2 = 400$ ;  $\therefore PQ=20$ .


7. By art. 20,  $PQ^2 = (-12-16)^2 + (15-36)^2 = 28^2 + 21^2 = 1225$ ;  $\therefore PQ=35$ .

8. Here  $y$  may be taken as the common ordinate of the point of intersection. From the 1st and 2nd equations therefore we obtain  $x = -16$ ,  $y = -15$ ; which values being put for  $x$  and  $y$  in the 3rd equation form the identity  $-45 = -32 - 13$ . Therefore the three lines meet at one point.

9. Taking  $y$  to denote the common ordinate of the point of intersection, we have  $\frac{1}{2}x + 3 = 6x - 12$ ; whence we obtain  $x = \frac{30}{11}$ ,  $y = \frac{48}{11}$ ; and, by substitution,  $\frac{48}{11} = -\frac{3}{11}m + 8$ ; which gives  $m = \frac{4}{3}$ .

10. From the given equations we obtain  $x = -5$ ,  $y = 4$ , which values being made the coordinates of a point to which a line is drawn from the origin, the equation to that line will be  $4x = -5y$ , or  $4x + 5y = 0$ .

11. Take  $OM$ ,  $AM$ ,  $= 21$ ,  $25$ ;  $ON$ ,  $BN$ ,  $= 5$ ,  $26$ ;  $OC = 46$ ;



and join  $AB$ ,  $BC$ ,  $CA$ , to form the proposed triangle.

By similar triangles we have

$$\frac{ME}{EN} = \frac{AM}{BN}; \text{ or } \frac{ME}{MN} = \frac{AM}{AM+BN}, \text{ that is, } \frac{ME}{26} = \frac{25}{51};$$

$$\text{hence } ME = \frac{650}{51}, \text{ and } EC = 67 - ME = \frac{2767}{51}.$$

The required area is  $= \frac{1}{2}EC(AM+BN) = 2767 \div 2 = 1383\frac{1}{2}$ .

12. Take, as the coordinates of the three points, OL, AL, =17, 20; OM, BM, =44, 33; ON, CN, =51, 8.

The triangle is = the trapezium ALMB + the trap<sup>m</sup> BMNC—the trap<sup>m</sup> ALNC;

$$= \frac{1}{2}(AL + BM)LM + \frac{1}{2}(BM + CN)MN - \frac{1}{2}(AL + CN)LN;$$


$$= \frac{1}{2}(53 \times 27 + 41 \times 7 - 28 \times 34) = 383.$$

13. (i.) Let the coordinates of P be  $x', y'$ , and let those of Q be  $x'', y''$ . The area of OPQ is

$$= \text{area of OPM} + \text{area of PMNQ} - \text{area of OQN};$$

$$= \frac{1}{2}x'y' + \frac{1}{2}(x'' - x')(y'' + y') - \frac{1}{2}x''y'';$$

$$= \frac{1}{2}(x'y' - x'y'').$$



(ii.)  $OP^2 = OM^2 + PM^2 = 25 + 144$   
 $= 169;$

$$OQ^2 = ON^2 + QN^2 = 144 + 81 = 225;$$

$$PQ^2 = (PM - QN)^2 + MN^2 = (12 - 9)^2 + (12 - 5)^2 = 58;$$

$$\therefore OP, OQ, PQ, = 13, 15, \sqrt{58}.$$

14. From 1st and 3rd equations we obtain  $x=2, y=3$ ;


from 1st and 2nd we have

$$x=5, y=2;$$

from 2nd and 3rd,

$$x=-8, y=28.$$

These coordinates denote the points P, Q, R, respectively.



$$\text{Area of RLQ} = \frac{1}{2}(28 + 2) \times 13 = 195$$

$$\text{do. RLMP} = \frac{1}{2}(28 + 3) \times 10 = 155$$

$$\text{do. PMNQ} = \frac{1}{2}(3 + 2) \times 3 = 7\frac{1}{2}$$

$$\underline{\quad 162\frac{1}{2}}\quad$$

$$\text{Area of PQR} = 32\frac{1}{2}$$

15. Take, as coordinates of the angular points, OL, AL, =3, 4; OM, BM, =8, 9; ON, CN, =10, 6. Also, for the coordinates of the point D take OH, DH, = $\frac{1}{2}(OL + OM)$

and  $\frac{1}{2}(AL+BM)$ ; and for those of E take  $OK, EK,$   
 $=\frac{1}{2}(OM+ON)$  and  $\frac{1}{2}(BM+CN)$ .

Thus,  $OH=\frac{1}{2}(3+8)=5\frac{1}{2}$ , and  $DH=\frac{1}{2}(4+9)=6\frac{1}{2}$ ;

also,  $OK=\frac{1}{2}(8+10)=9$ , and  $EK=\frac{1}{2}(9+6)=7\frac{1}{2}$ .

By art. 15, the equation to DE, passing through  $(5\frac{1}{2}, 6\frac{1}{2})$   
 and  $(9, 7\frac{1}{2})$ , is

$$y-6\frac{1}{2}=\frac{7\frac{1}{2}-6\frac{1}{2}}{9-5\frac{1}{2}}(x-5\frac{1}{2}); \text{ or } 14y=4x+69;$$

$$\text{whence } y=\frac{2}{7}x+\frac{6\frac{1}{2}}{7}.$$

The equation to AC, passing through  $(3, 4)$  and  $(10, 6)$ , is

$$y-4=\frac{6-4}{10-3}(x-3); \text{ or } 7y=2x+22;$$

$$\text{whence } y=\frac{2}{7}x+\frac{22}{7}.$$

Hence, the tangents of the angles which DE and AC make  
 with the axis of  $x$  being each  $=\frac{2}{7}$ , the lines DE and AC are  
 parallel.

16. Take  $ON=a, OM=a', OL=b, OK=b'$ . DN is the  
 line  $x=a$ , CM is  $x=a'$ .

LB is  $y=b$ , KC is  $y=b'$ .

The equation to AC, through  
 $(a, b)$  and  $(a', b')$ , is

$$y-b=\frac{b'-b}{a'-a}(x-a);$$

$$\text{or, } (a'-a)y-(b'-b)x-a'b$$


$$+ab=0.$$

The equation to DB, through  $(a, b')$  and  $(a', b)$ , is

$$y-b'=\frac{b-b'}{a'-a}(x-a); \text{ or, } (a'-a)y+(b'-b)x-a'b'+ab=0.$$

17. In OX take the abscissa  $OM=12$ , draw the ordinate  
 $PM=\frac{3}{4}$  of  $12=9$ , and produce MP to Q making  $QM=\frac{3}{4}$  of  
 $12=23$ . Join OP, OQ, and complete the parallelogram  
 OPQR.

PQ parallel to the axis of  $y$  is the line  $x=12$ ; OP, whose  
 coordinates are 12, 9, is the line  $4y=3x$ ; and OQ, whose  
 coordinates are 12, 23, is the line  $12y=23x$ . The equation





to OR on the axis of  $y$  is  $x=0$ ; also  $OR=PQ=QM-PM=14$ .

The equation to QR, through  $(0, 14)$  and  $(12, 23)$ , is

$$y-14=\frac{23-14}{12}x; \text{ or } 4y-3x=56.$$

The equation to PR, through  $(0, 14)$  and  $(12, 9)$ , is

$$y-14=\frac{9-14}{12}x; \text{ or } 5x+12y=168.$$

The area of the parallelogram  $=2^{\text{nd}}$  the difference of the triangles OQM, OPM,  $=OM(QM-PM)=12(23-9)=168$ .

18. The equation to the diagonal AC is

$$y-3=\frac{5-3}{3-2}(x-2); \text{ or } y=2x-1.$$

We have now to find the equations to the lines AB, AD, passing through  $(2, 3)$ , and making an angle of  $45^\circ$  with the line  $y=2x-1$ . In art. 19 we have the applicable formula

$$y-y'=\frac{m\pm\tan\phi}{1\mp m\tan\phi}(x-x');$$

$$\therefore y-3=\frac{2+1}{1-2}(x-2); \text{ or } y=-3x+9;$$

$$\text{and } y-3=\frac{2-1}{1+2}(x-2); \text{ or } 3y=x+7;$$

these are the equations to AD and AB.

Also, we have to find the equations to CB, CD, passing through  $(3, 5)$ , and making an angle of  $45^\circ$  with the line  $y=2x-1$ .

$$y-5=\frac{2+1}{1-2}(x-3); \text{ or } y=-3x+14;$$

$$y-5=\frac{2-1}{1+2}(x-3); \text{ or } 3y=x+12;$$

these are the equations to CB and CD.

19. The equation to OP is  $y=\frac{1}{2}x$ ; and that to PQ is  $y=-\frac{7}{4}x+7$ . Now, since every point on the bisecting line

is equidistant from OP and PQ, let  $(x, y)$  be any point on the bisecting line; its perpendicular distance from OP, by art. 21, is

$$\frac{y - \frac{1}{2}x - 0}{\sqrt{1 + \frac{1}{4}}} ; \text{ or } \frac{5y - 12x}{13} ;$$

also its perpendicular distance from PQ is

$$\frac{y + \frac{7}{4}x - 7}{\sqrt{1 + \frac{49}{16}}} ; \text{ or } \frac{24y + 7x - 168}{25} ;$$

$$\therefore \frac{5y - 12x}{13} = \frac{24y + 7x - 168}{25} ; \text{ or } 391x + 187y = 2184.$$

20. Take A as the origin, AB, AC as the axes. Let  $AB = a, AC = b$ .

Because BC cuts off intercepts  $a$  and  $b$  on the axes, its equation is

$$\frac{x}{a} + \frac{y}{b} = 1 ; \text{ or } y = -\frac{b}{a}x + b. \quad (1)$$

Again, since AK and AF are squares, P is the point  $(-b, -a)$ ; and the equation to PA passing through the

origin is

$$y = \frac{a}{b}x. \quad (2)$$

Accordingly (1) and (2), having reciprocal coefficients of  $x$  with opposite signs, are perpendicular to each other.

21. Here the given equations supply the respective tangents of CAX and CBX, viz.  $m'$  and  $m$ ;  $\therefore \tan CBA = -m$ .



$\tan ACB = \tan (CBX - CAX)$

$$= \frac{m - m'}{1 + mm'}.$$

The sides may be found as follows:

AD being  $= c$ , and  $\frac{AB}{AD} = \frac{BR}{CR}$

$= -\frac{1}{m}$ , we have the magnitude of  $AB = \frac{c}{m}$ .



Then  $\frac{CR}{AR} = m'$ ;  $\frac{BR}{CR} = -\frac{1}{m}$ ;  $\therefore \frac{AR}{BR} = \frac{m}{-m'}$ ;  $\frac{AR}{AB} = \frac{m}{m-m'}$ ;

hence,  $AR = \frac{c}{m-m'}$ ,  $CR = \frac{cm'}{m-m'}$ ,  $BR = \frac{cm'}{m(m-m')}$ .

$$AC^2 = AR^2 + CR^2 = \frac{c^2}{(m-m')^2} (1+m'^2);$$

$$\text{or } AC = \frac{c}{m-m'} \sqrt{1+m'^2};$$

$$BC^2 = BR^2 + CR^2 = \frac{c^2 m'^2}{(m-m')^2} \cdot \frac{1+m^2}{m^2};$$

$$\text{or } BC = \frac{cm'}{m(m-m')} \sqrt{1+m^2}.$$

### EXERCISES [C].

1. (i.) Completing squares with  $x^2-6x$  and  $y^2-8y$ , we have

$$(x-3)^2 + (y-4)^2 = 24 + 9 + 16 = 49.$$

Therefore the radius  $= \sqrt{49} = 7$ ; and the coordinates of the centre are 3 and 4.

(ii.) Completing squares with  $x^2+2x$  and  $y^2+4y$ , we have

$$(x+1)^2 + (y+2)^2 = 4 + 1 + 4 = 9.$$

Therefore the radius  $= \sqrt{9} = 3$ ; and the coordinates of the centre are  $-1$  and  $-2$ .

(iii.) Completing squares with  $x^2-6x$  and  $y^2-18y$ , we have

$$(x-3)^2 + (y-9)^2 = 80 = 16 \times 5.$$

Therefore the radius  $= 4\sqrt{5}$ ; and the coordinates of the centre are 3 and 9.

(iv.) Completing squares with  $x^2+x$  and  $y^2-2y$ , we have

$$(x+\frac{1}{2})^2 + (y-1)^2 = 5.$$

Therefore the radius  $= \sqrt{5}$ ; and the coordinates of the centre  $-\frac{1}{2}$  and 1.

2. (i.) The second equation gives  $y^2 = x^2 - 4x + 4$ ; therefore  $x^2 - 4x + 4 = 34 - x^2$ ; whence  $x = 5$  or  $-3$ ;  $y = 3$  or  $-5$ ; so that the points of intersection are  $(5, 3)$  and  $(-3, -5)$ .

(ii.) The 2nd equation gives  $y^2 = 4x^2 + 4x + 1$ ; therefore  $4x^2 + 4x + 1 = 34 - x^2$ ; whence  $x = 2\frac{1}{2}$  or  $-3$ ;  $y = -5\frac{1}{2}$  or  $5$ ; so that the points of intersection are  $(2\frac{1}{2}, -5\frac{1}{2})$  and  $(-3, 5)$ .

3. (i.) The 2nd equation gives  $16y^2 = 9x^2 - 30x + 25$ ; therefore  $9x^2 - 30x + 25 = 16 - 16x^2$ ; or  $25x^2 - 30x + 9 = 0$ ;  $\therefore 5x - 3 = 0$ ; whence  $x = \frac{3}{5}$ ;  $y = \frac{1}{4}(3x - 5) = -\frac{4}{5}$ ; so that the line meets the circle at only one point, being a tangent to it at the point  $(\frac{3}{5}, -\frac{4}{5})$ .

(ii.) The second equation gives  $16y^2 = 25x^2 + 50x + 25$ ; therefore  $25x^2 + 50x + 25 = 16 - 16x^2$ ; whence  $x = -1$  or  $-\frac{9}{4}$ ;  $y = -\frac{1}{4}(5x + 5) = 0$  or  $-\frac{4\frac{1}{2}}{1}$ ; so that the line meets the circle at the points  $(-1, 0)$  and  $(-\frac{9}{4}, -\frac{4\frac{1}{2}}{1})$ , the former of these points being an extremity of the diameter in the axis of  $x$ .

4. (i.) From the two given equations we obtain

$$16y^2 = 9x^2 - 192x + 1024 = -16x^2 + 384x + 160y;$$

whence, substituting for  $160y$  its value in terms of  $x$ , viz.

$$120x - 1280, \text{ we have } 25x^2 - 696x = -2304;$$

$\therefore x = 24$  or  $\frac{32}{5}$ ;  $y = \frac{1}{4}(3x - 32) = 10$  or  $-\frac{1\frac{1}{2}}{5}$ ; so that the points of intersection (the origin being on the circumference) are  $(24, 10)$  and  $(3\cdot84, -5\cdot12)$ , the former of these being an extremity of the diameter through the origin.

(ii.) From the 1st equation  $y = \frac{3}{4}x - 8$ , and  $2y^2 = 1\frac{1}{8}x^2 - 24x + 128$ ; hence, substituting these values in the 2nd equation, we have  $x^2 - 10x = -24$ ;  $\therefore x = 6$  or  $4$ ,  $y = -3\frac{1}{2}$  or  $-5$ ; so that the points of intersection are  $(6, -3\frac{1}{2})$  and  $(4, -5)$ .

5. Let  $(x', y')$  be the point on the circle at which the tangent is drawn: The equation to the tangent (see art 28) is

$$(x-a)(x'-a) + (y-b)(y'-b) = c^2;$$

$$\text{which gives } y-b = -\frac{x'-a}{y'-b}(x-a) + \frac{c^2}{y'-b};$$

or, expressing  $\frac{c^2}{y'-b}$  in terms of the tangent  $-\frac{x'-a}{y'-b}$ , we have, by art. 29,

$$y-b = -\frac{x'-a}{y'-b}(x-a) \pm c\sqrt{\frac{(x'-a)^2}{(y'-b)^2} + 1}.$$

Now, in order that this line may be parallel to the line  $y=mx+n$ , we must have  $-\frac{x'-a}{y'-b}=m$ ; hence the required equation is

$$y-b = m(x-a) \pm c\sqrt{1+m^2}.$$

6. The equation to the circle may be written

$$(x-20)^2 + y^2 = 400;$$

so that the radius is 20. The equation to the straight line, AQ, may be written

$$\frac{x}{-11\frac{1}{2}} + \frac{y}{4\frac{2}{3}} = 1; \text{ so that } OA = 11\frac{1}{2}, OK = 4\frac{2}{3};$$

$$\therefore AK = \sqrt{(OA^2 + OK^2)} = 18\frac{2}{3}.$$

Now, by art. 31, the perpendicular CR from the centre bisects the chord PQ; and by similar triangles we have

$$\frac{CR}{AC} = \frac{OK}{AK}, \text{ or } CR = \frac{1}{3} \times \frac{15\frac{1}{2}}{18\frac{2}{3}} \times \frac{15\frac{1}{2}}{18\frac{2}{3}} = 12;$$


hence  $PR = \sqrt{(OC^2 - CR^2)} = 16$ ;  $\therefore PQ = 32$ .

7. Squaring the equation to the line gives

$y^2 = \frac{b^2}{a^2}x^2 - \frac{2b^2}{a}x + b^2$ ; whence by substitution in the equation to the circle we have

$$\frac{a^2 + b^2}{a^2}x^2 - \frac{2b^2}{a}x + b^2 - c^2 = 0,$$

$$\text{or, } x^2 - \frac{2ab^2}{a^2 + b^2}x + \frac{a^2(b^2 - c^2)}{a^2 + b^2} = 0.$$



This quadratic will have two roots, except when the third term is the square of half the coefficient of the second term; that is, when

$$\frac{a^2(b^2 - c^2)}{a^2 + b^2} = \left( \frac{ab^2}{a^2 + b^2} \right)^2; \text{ or when } (b^2 - c^2)(a^2 + b^2) = b^4;$$

$$\text{or when } b^2c^2 + a^2c^2 = a^2b^2;$$

which, by dividing each term by  $a^2b^2c^2$ , becomes


$$\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{c^2}.$$

8. Let O, the centre of the circle whose radius is  $c$ , be the origin of coordinates: The equation to that circle is  $x^2 + y^2 = c^2$ .

Now  $CR^2 + QR^2 = CQ^2$ , or  $(a - x)^2 + y^2 = c'^2$ ; or  $a^2 - 2ax + x^2 + y^2 = c'^2$ ;

that is,  $a^2 - 2ax + c^2 = c'^2$ .

$$\therefore x = \frac{1}{2a} (a^2 + c^2 - c'^2).$$



$$\text{Again; } y^2 = c^2 - x^2 = c^2 - \frac{1}{4a^2} (a^2 + c^2 - c'^2)^2;$$

$$\begin{aligned} \therefore y &= \pm \frac{1}{2a} \sqrt{4a^2c^2 - (a^2 + c^2 - c'^2)^2} \\ &= \pm \frac{1}{2a} \sqrt{(2ac + a^2 + c^2 - c'^2)(2ac - a^2 - c^2 + c'^2)} \\ &= \pm \frac{1}{2a} \sqrt{\{(a+c)^2 - c'^2\} \{c'^2 - (a-c)^2\}} \end{aligned}$$

$$\text{or } y = \pm \frac{1}{2a} \sqrt{(a+c+c')(a+c-c')(a-c+c')(c-a+c')}.$$

9. The general equation to the circle being

$$x^2 + y^2 + Ax + By + C = 0,$$

and B being in the present case = 0, the centre is on the axis of  $x$  at the distance  $\frac{1}{2}A = \frac{m^2 + 1}{m^2 - 1}a$  from the origin. The

$$\text{radius} = \sqrt{\frac{1}{4}A^2 - a^2} = a \sqrt{\left( \frac{m^2 + 1}{m^2 - 1} \right)^2 - 1}$$

$$= a \sqrt{\frac{2m^2}{m^2 - 1} \cdot \frac{2}{m^2 - 1}} = \frac{2m}{m^2 - 1}a.$$

To determine the points at which the circle cuts the axis of  $x$ , make  $y=0$  in the general equation, and the values of  $x$  will then be found

$$= -\frac{1}{2}A \pm \sqrt{\frac{1}{4}A^2 - C}; = \frac{m^2+1}{m^2-1}a \pm \frac{2m}{m^2-1}a,$$

$$= \frac{m^2 \pm 2m + 1}{m^2 - 1}a, = \frac{m+1}{m-1}a \text{ and } \frac{m-1}{m+1}a.$$

10. Take O, the middle point of the base AB, as origin, AO=OB=a; and let  $(x, y)$  be the vertex C.

$$AC^2 = CM^2 + AM^2 = y^2 + (a+x)^2;$$

$$BC^2 = CM^2 + BM^2 = y^2 + (a-x)^2;$$

whence, by addition,

$$2(y^2 + x^2 + a^2) = AC^2 + BC^2,$$

that is,

$$2(OC^2 + OA^2) = AC^2 + BC^2.$$

Again; putting  $s$  for  $AC^2 + BC^2$ ,

$$y^2 + x^2 + a^2 = \frac{1}{2}s; \text{ or } x^2 + y^2 = \frac{1}{2}s - a^2,$$

the equation to a circle whose radius  $= \sqrt{\frac{1}{2}s - a^2}$ .

11. Take the intercept OB=5, OT= $\frac{1}{3}$  of 5. TB is the line touching the required circle.

$$TB^2 = OB^2 + OT^2 = 2\frac{5}{3}^2;$$

$$\therefore TB = \frac{1}{3}\sqrt{250}.$$



Now, OP·TB=OB·OT, that is,

$$OP \times \frac{1}{3}\sqrt{250} = 5 \times \frac{5}{3};$$

$\therefore OP^2 = 625 \div 250 = 2\frac{1}{2}$ ; and the equation to the circle is, therefore,

$$x^2 + y^2 = 2\frac{1}{2}.$$

12. From the given equations we obtain  $x = \frac{1}{2}a$ , and  $y = \frac{1}{2}\sqrt{4c^2 - a^2}$ ; that is, the coordinates of the centre of the required circle are  $\frac{1}{2}a$  and 0; and those of one of the points of intersection of the two given circles are  $\frac{1}{2}a$  and  $\frac{1}{2}\sqrt{4c^2 - a^2}$ ; therefore, the square of the radius of the required circle is  $(\frac{1}{2}\sqrt{4c^2 - a^2})^2 = c^2 - \frac{1}{4}a^2$ .



Hence the required equation is

$$(x - \frac{1}{2}a)^2 + (y - 0)^2 = c^2 - \frac{1}{4}a^2; \text{ or, } x^2 + y^2 - ax = c^2 - \frac{1}{2}a^2.$$

13. The two centres, P and Q, are the points  $(a, b)$  and  $(b, a)$ , and, by art. 20, the length of the line joining these points is determined by

$$PQ^2 = (b - a)^2 + (a - b)^2 = 2(a - b)^2;$$

hence the distance between the centres is  $\sqrt{2(a - b)^2}$ ; and as the radii are equal, we have  $c = \pm \frac{1}{2}\sqrt{2(a - b)}$ .

14. The equation  $x^2 + y^2 - 2cx = 0$  denotes that the origin is the extremity of a diameter in the axis of  $x$ .

The condition of intersection of the circle with the line  $y = mx$ , or  $y^2 = m^2x^2$ , is  $2cx - x^2 = m^2x^2$ ; whence  $x = \frac{2c}{1 + m^2}$ ,  $y = \frac{2mc}{1 + m^2}$ ; and the square of the line  $= \frac{4c^2 + 4m^2c^2}{(1 + m^2)^2} = \frac{4c^2}{1 + m^2}$ ;  $\therefore$  square of radius of required circle  $= \frac{c^2}{1 + m^2}$ .

The coordinates, then, of the centre of the required circle being  $\frac{c}{1 + m^2}$  and  $\frac{mc}{1 + m^2}$ , we have, for the required equation,

$$\begin{aligned} \left(x - \frac{c}{1 + m^2}\right)^2 + \left(y - \frac{mc}{1 + m^2}\right)^2 &= \frac{c^2}{1 + m^2}; \\ \text{or } x^2 + y^2 - \frac{2c}{1 + m^2}x - \frac{2mc}{1 + m^2}y + \frac{c^2}{1 + m^2} &= \frac{c^2}{1 + m^2}; \\ \text{that is, } x^2 + y^2 - \frac{2c}{1 + m^2}(x - my) &= 0. \end{aligned}$$

#### EXERCISES [D].

1. (i.) Dividing by  $c^2$ , we have

$$\frac{x^2}{\frac{1}{3}c^2} + \frac{y^2}{\frac{1}{4}c^2} = 1; \therefore a^2 = \frac{1}{3}c^2, \text{ and } b^2 = \frac{1}{4}c^2;$$

$$\therefore \frac{b^2}{a^2} = \frac{3}{4}; \text{ and } 1 - \frac{b^2}{a^2} = e^2 = \frac{1}{4}; \therefore e = \frac{1}{2}.$$



(ii.) Dividing by  $c'^2$ , we have

$$\frac{x^2}{\frac{1}{3}c'^2} + \frac{y^2}{\frac{1}{3}c'^2} = 1; \therefore a^2 = \frac{1}{3}c'^2, \text{ and } b^2 = \frac{1}{3}c'^2;$$

$$\therefore \frac{b^2}{a^2} = \frac{3}{5}; \text{ and } 1 - \frac{b^2}{a^2} = e^2 = \frac{2}{5} = \frac{1}{2}\frac{2}{5}; \therefore e = \frac{1}{2}\sqrt{10}.$$

(iii.) Dividing by  $780^2$  shews  $a^2 = 156^2$ , and  $b^2 = 60^2$ ;

$$\therefore \frac{b^2}{a^2} = \frac{25}{169}; \text{ and } 1 - \frac{b^2}{a^2} = e^2 = \frac{144}{169}; \therefore e = \frac{12}{13}.$$

2. The coordinates of the points A' and B are  $-a, 0$ , and  $0, b$ ; those of C and F are  $0, 0$ , and  $ae, \frac{b^2}{a}$ ; hence the equations to the lines A'B and CF are, respectively,

$$y = \frac{b}{a}x + b, \text{ and } y = \frac{b^2}{a^2} \cdot \frac{1}{e}x;$$

and that these lines may be parallel they must have the same inclination to the axis of  $x$ , or the tangents  $\frac{b}{a}$  and  $\frac{b^2}{a^2} \cdot \frac{1}{e}$  must be equal;

that is,  $\sqrt{1-e^2} = \frac{1}{e}(1-e^2)$ , or  $1 = \frac{1}{e^2}(1-e^2)$ ;  $\therefore e = \frac{1}{2}\sqrt{2}$ .

3.  $a = \frac{1}{2}(4+6) = 5$ ;  $SC = ae = 4$ ;  $SH = 8$ ;

$SM^2 - MH^2 = SP^2 - HP^2$ ; whence  $SM - MH = 2\frac{1}{2}$ ;

$\therefore SM = 5\frac{1}{4}$ ;  $x' = CM = SM - SC = 1\frac{1}{4}$ .

$PM^2 = (a^2 - x'^2)(1 - e^2) = \frac{9}{16}$  of 15;  $\therefore y' = PM = \frac{3}{4}\sqrt{15}$ .

4. The equation to the normal at the point  $(x', y')$  is

$$y - y' = \frac{a^2 y'}{b^2 x'}(x - x').$$

In the present case, we have  $x' = CH = ae$ , and  $y' = HF = \frac{b^2}{a}$ ; and therefore  $\frac{a^2 y'}{b^2 x'} = \frac{ab^3}{ab^2 e} = \frac{1}{e}$ ; hence the equation becomes

$$y - \frac{b^2}{a} = \frac{1}{e} (x - ae) = \frac{x}{e} - a,$$

$$\text{or } y + \frac{a^2 - b^2}{a} = \frac{x}{e};$$

but  $b^2 = a^2 - a^2e^2$ , or  $a^2 - b^2 = a^2e^2$ ; therefore, the equation to the normal at F is  $y + ae^2 = \frac{x}{e}$ .

$$5. y'^2 = (a^2 - x'^2)(1 - e^2);$$

$$\text{that is, } \frac{49 \times 119}{32} = (a^2 - \frac{7^2}{32}) \times \frac{4}{81};$$

$$\text{or, } 119 = (32a^2 - 729) \times \frac{1}{81}; \text{ or } a^2 = 324; \therefore a = 18.$$

$$SP = a + ex = 18 + \frac{4}{3} \sqrt{2} \times \frac{7}{8} \sqrt{2} = 18 + 3 = 21;$$

$$HP = a - ex = 18 - 3 = 15.$$

6. The equation  $2x^2 + 3y^2 = 18$  may be written

$$\frac{x^2}{9} + \frac{y^2}{6} = 1, \text{ where } 9 = a^2, 6 = b^2.$$

Now, from art. 46, the equation to the tangent is

$$y = mx \pm \sqrt{a^2m^2 + b^2};$$

and we have here  $m = \tan 30^\circ = \frac{1}{\sqrt{3}}$ ;


$$\therefore y = \frac{1}{\sqrt{3}} \sqrt{3}x \pm \sqrt{3 + 6}; \text{ or } 3y = x\sqrt{3} \pm 9.$$

7. Given  $a^2 = 2b^2$ ;  $\therefore a^2 - b^2 = b^2 = \frac{1}{2}a^2$ ; that is,  $a^2e^2 = \frac{1}{2}a^2$ ;

$$\therefore 4a^2e^2 = 2a^2; \text{ that is, } SH^2 = SB^2 + HB^2.$$

8. (See fig. in art. 49.) Since  $\sin VCT = \cos VTC = \cos \phi$ , we have, from art. 51, the perpendicular to the tangent, from the centre as origin,  $= a(1 - e^2 \cos^2 \phi)^{\frac{1}{2}}$ ; and it is an arithmetical mean between HR and SR'. Suppose CZ drawn perpendicular to SR', then  $SZ = SC \sin \angle SCZ = ae \sin \phi$ ;  $\therefore SR'$  and  $HR = a\{e \sin \phi \pm (1 - e^2 \cos^2 \phi)^{\frac{1}{2}}\}$ .

9. Let BC, PM, QN be ordinates of an ellipse, and let them be produced to B', P', Q', points in the circumference of a circle described on the transverse axis.



By art. 49 we have

$$\frac{BC}{B'C} = \frac{PM}{P'M} = \frac{QN}{Q'N} = \frac{b}{a};$$

$$\therefore \frac{BC+PM}{B'C+P'M} = \frac{PM+QN}{P'M+Q'N} = \frac{b}{a}.$$

Now, the areas of the trapeziums BCMP and B'CMP' are  $\frac{1}{2}CM(BC+PM)$  and  $\frac{1}{2}CM(B'C+P'M)$ , and are therefore as  $b$  to  $a$ ; and the same ratio holds for the trapeziums PN and P'N, and for every pair of trapeziums similarly situated; so that a polygon of an indefinite number of sides inscribed in the ellipse will be to the corresponding polygon in the circle as  $b$  to  $a$ ; that is, the area of the ellipse is to that of the circle as  $b$  to  $a$ .

Accordingly, since the area of the circle described on the transverse axis of the ellipse is  $\pi a^2$ , we have

$$a : b :: \pi a^2 : \pi ab, \text{ the area of the ellipse.}$$

Hence, if  $r$  be the radius of a circle equal in area to an ellipse whose semi-axes are  $a$  and  $b$ , we have  $\pi ab = \pi r^2$ ; that is,  $ab = r^2$ , or the radius is a mean proportional between the semi-axes.

10. In the given ellipse put  $c^2$  for  $a^2$ , as the  $a$  here has not the same import as that in the general equation; then  $a^2 = 9c^2$ , and  $b^2 = 4c^2$ ;  $\therefore a^2 - b^2 = a^2 e^2$  becomes  $5c^2 = 9e^2 c^2$ ; hence  $e = \frac{1}{3}\sqrt{5}$ .

The equation to the tangent at the point  $(x', y')$  is

$$y = -\frac{b^2 x'}{a^2 y'} x + \frac{b^2}{y'}; \text{ which, for } x' = ae, y' = \frac{b^2}{a}, \text{ becomes}$$

$y = -ex + a$ . Now, instead of  $a$  resume  $\sqrt{9a^2}$ , or  $3a$ , from the given form of equation, and the required equation will be


$$y = -\frac{1}{3}\sqrt{5}x + 3a;$$

$$\text{or, } 3y + \sqrt{5}x - 9a = 0.$$

From the first of these forms we see that the required intercept on the axis of  $y$  is  $3a$ ; and that on the axis of  $x$  is

$$3a + \frac{1}{3}\sqrt{5} = 9a + \sqrt{5} = \frac{9a}{5}\sqrt{5}.$$

11. Let S and H be the centres of the given circles, P the centre of an intervening circle touching the given circles at A and B.



Join SP, HP, which will pass through the points of contact.

$$HP + SP = HA + AP + SB - BP; \text{ but } AP = BP;$$

$\therefore HP + SP = HA + SB$ , the sum of the given radii, and therefore constant.

And the locus of a point, the sum of whose distances from two given points is constant, is an ellipse. (See the 4th of the worked Examples.)

12. If  $x', y'$  were the coordinates of the point P: the equation to A'P passing through  $(-a, 0)$  and  $(x', y')$  would be

$$y = \frac{y'}{x' + a} (x + a);$$

and the equation to AP through  $(a, 0)$  and  $(x', y')$  would be

$$y = \frac{y'}{x' - a} (x - a).$$

Hence, by art. 17, the tangent of the angle of intersection of A'P and AP is

$$= \left( \frac{y}{x - a} - \frac{y}{x + a} \right) \div \left( 1 + \frac{y^2}{x^2 - a^2} \right) = \frac{2ay}{x^2 - a^2 + y^2}.$$

But, as the lines are to intersect on the ellipse, we have

$$a^2 y^2 + b^2 x^2 = a^2 b^2; \text{ or, } x^2 - a^2 = -\frac{a^2}{b^2} y^2;$$

whence, by substitution, the tangent of the angle of intersection becomes  $= 2ay \div \left( 1 - \frac{a^2}{b^2} \right) y^2$

$$= \frac{-2ab^2}{(a^2 - b^2)y} = \frac{-2ab^2}{a^2 e^2 y} = \frac{-2b^2}{ae^2 y}.$$

## EXERCISES [E].

1. Here  $e^2 - 1 = 4 - 1 = 3 = \frac{b^2}{a^2}$ ;  $\therefore b^2 = 3a^2$ ; and the equation to the hyperbola, viz.  $y^2 = \frac{b^2}{a^2}x^2 - b^2$ , becomes

$$y^2 = 3(x^2 - a^2).$$

2. Here we have  $3x^2 - 2y^2 = -6n^2$ ; or,  $\frac{x^2}{-2n^2} - \frac{y^2}{-3n^2} = 1$ ;

$$\therefore a^2 : b^2 :: 2n^2 : 3n^2 :: 2 : 3;$$

$$e^2 - 1 = \frac{b^2}{a^2} = \frac{3}{2}; \text{ or } e^2 = \frac{5}{2} = \frac{10}{4}; \therefore e = \frac{1}{2}\sqrt{10}.$$

And, since  $\frac{b^2}{a^2} = \frac{3}{2}$ ,  $\therefore \frac{2b^2}{a} = 3a$ , the latus rectum.

3. In the ellipse the coordinates of A are  $x' = a$ ,  $y' = 0$ ; for that point, therefore, the equation to the tangent,

$$\text{viz. } a^2yy' + b^2xx' = a^2b^2,$$

$$\text{becomes } x = a,$$

which is the equation to a line through A parallel to CY.

In the hyperbola, the coordinates of A are  $x' = a$ ,  $y' = 0$ ; for that point, therefore, the equation to the tangent,

$$\text{viz. } a^2yy' - b^2xx' = -a^2b^2,$$

$$\text{again becomes } x = a.$$

4. The equation to the tangent at P is

$$a^2yy' - b^2xx' = -a^2b^2.$$

The intercept CT, made by the tangent TP, is found from the equation to the tangent by putting  $y = 0$ ; whence

$$x = \frac{a^2}{x'} = CT; \therefore \frac{CT}{CA} = \frac{a}{x'} = \frac{CA}{CM} = \frac{CE}{CP};$$

$\therefore$  CTE and CAP are similar triangles.

5. From  $\frac{y}{m} - \frac{x}{n} = 1$  we have  $y = \frac{m}{n}x + m$ ; (1)

and the tangent to the hyperbola being of the form

$$y = m'x \pm \sqrt{a^2m'^2 - b^2}, \quad (2)$$

$m'$ , the tangent of inclination to the transverse axis, is  $= \frac{m}{n}$ ;  $\therefore$  (2) becomes

$$y = \frac{m}{n}x \pm \sqrt{a^2 \frac{m^2}{n^2} - b^2};$$

hence, 
$$\frac{m}{n}x + m = \frac{m}{n}x \pm \sqrt{a^2 \frac{m^2}{n^2} - b^2},$$

or,  $a^2 \frac{m^2}{n^2} - b^2 = m^2$ ; or,  $\frac{a^2}{n^2} - \frac{b^2}{m^2} = 1$ .

6. The equation to a line touching the ellipse (*see art. 46*) is  $y = mx + \sqrt{a^2 m^2 + b^2}$ ; (1)

that to a line touching the hyperbola is of the form

$$y = m'x + \sqrt{a'^2 m'^2 - b'^2}. \quad (2)$$

In order that (2) may be at right angles to (1) we must have  $m' = -\frac{1}{m}$ ; hence (2) becomes

$$y = -\frac{1}{m}x + \frac{\sqrt{a'^2 - b'^2 m^2}}{m},$$

$$\text{or } my = -x + \sqrt{a'^2 - b'^2 m^2}. \quad (3)$$

Squaring (1) and (3) and then adding the results, we have

$$1 + m^2(x^2 + y^2) = a'^2 + b^2 + (a^2 - b'^2)m^2; \quad (4)$$

but, since the curves have the same foci and centre,

$$SC^2 = a^2 - b^2 = a'^2 + b'^2; \therefore a^2 - b'^2 = a'^2 + b^2.$$

Thus (4) becomes


$$(1 + m^2)(x^2 + y^2) = (a'^2 + b^2)(1 + m^2),$$

$$\text{or, } x^2 + y^2 = a'^2 + b^2;$$

which is the equation to a circle passing through the four points of intersection of the curves.


7. If N, the fixed point in AB, is between A and B, and  $AN > NB$ , we have

$AP - PB = AD - BE = AN - NB$ , a constant difference; hence, in this case, the locus of P is an hyperbola of which A and B are the foci.



If, however,  $AN=NB$ , the locus will be the perpendicular to  $AB$  at  $N$ .

But if the point  $N$  be on  $AB$  produced, then  $AP + PB = AD + EB = AN + BN$ , a constant sum; so that in this case the locus of  $P$  is an ellipse.



$$8. \text{ Magnitude of } \cos PSH = \frac{SM}{SP} = \frac{x+ae}{ex+a};$$

$$\tan^2 \frac{1}{2}PSH = \frac{\sin^2 \frac{1}{2}PSH}{\cos^2 \frac{1}{2}PSH} = \frac{1 - \cos PSH}{1 + \cos PSH}$$

$$= \frac{1 - \frac{x+ae}{ex+a}}{1 + \frac{x+ae}{ex+a}} = \frac{(e-1)(x-a)}{(e+1)(x+a)}.$$

$$\text{Magnitude of } \cos PHS = \frac{HM}{HP} = \frac{x-ae}{ex-a};$$


$$\tan^2 \frac{1}{2}PHS = \frac{1 - \frac{x-ae}{ex-a}}{1 + \frac{x-ae}{ex-a}} = \frac{(e-1)(x+a)}{(e+1)(x-a)}.$$

$$\therefore \tan \frac{1}{2}PSH \tan \frac{1}{2}PHS = \frac{e-1}{e+1}.$$

9. Let the coordinates of  $Q$  be  $x$  and  $y$ ; let those of  $P$  be  $x'$  and  $y'$ .

By similar triangles,

$$\frac{QN}{AN} = \frac{P'M}{AM}, \text{ or } \frac{y}{x-a} = \frac{y'}{a-x'};$$



$$\frac{QN}{A'N} = \frac{PM}{A'M}, \text{ or } \frac{y}{x+a} = \frac{y'}{a+x'};$$

$\therefore$  by mult<sup>n</sup>  $\frac{y^2}{a^2-x^2} = \frac{y'^2}{a^2-x'^2}$ . But the equation to the

ellipse is  $y'^2 = \frac{b^2}{a^2} (a^2 - x'^2)$ , or  $\frac{y'^2}{a^2 - x'^2} = \frac{b^2}{a^2}$ ;

$\therefore \frac{y^2}{x^2 - a^2} = \frac{b^2}{a^2}$ , or,  $y^2 = \frac{b^2}{a^2} (x^2 - a^2)$ , the equation to an hyperbola having the same axes as the ellipse.

10. The centre of the hyperbola being origin, the equation to the circle is

$$(x - ae)^2 + (y - 0)^2 = b^2 = a^2(e^2 - 1). \quad (1)$$

The equation to the asymptotes is  $y = \pm \frac{b}{a} x$ ;

$$\therefore y^2 = \frac{b^2}{a^2} x^2 = (e^2 - 1)x^2;$$

hence, by substitution in (1), that the circle may meet the asymptotes,

$$(x - ae)^2 + (e^2 - 1)x^2 = a^2(e^2 - 1);$$

$$\text{or } e^2 x^2 - 2aex + a^2 = 0; \text{ or, } (ex - a)^2 = 0;$$

$$\therefore x = \frac{a}{e}.$$

Accordingly, the circle touches each of the asymptotes at a point of which the abscissa is  $\frac{a}{e}$ , that is, at the point where the directrix meets them.

#### EXERCISES [F].

1. Since  $LS = 2AS$ , the equation to  $AL$  is  $y = 2x$ . The equation to a tangent is  $yy' = 2a(x + x')$ ; and in the present case  $y' = 2a$ ,  $x = a$ ;  $\therefore$  the equation to the tangent at  $L$  is  $2ay = 2a(x + a)$ , or  $y = x + a$ ; hence, by art. 17, the angle of intersection of the lines


$$y = 2x \text{ and } y = x + a,$$

is that whose tangent is  $\frac{2-1}{1+2 \times 1}$ , or  $\frac{1}{3}$ .





7. Let  $x, y$  be the coordinates of  $P$ ,  $x', y'$  those of  $P'$ .



By similar triangles,  $\frac{SM}{SM'} = \frac{SP}{SP'} = \frac{DM}{DM'}$ ;  $\therefore \frac{SM}{DM} = \frac{SM'}{DM'}$ ,

that is,  $\frac{x-a}{x+a} = \frac{a-x'}{a+x'}$ ;  $\therefore \frac{a}{x} = \frac{x'}{a}$ . (1)

Now,  $y^2 = 4ax$ , and  $y'^2 = 4ax'$ ;  $\therefore x = \frac{y^2}{4a}$ , and  $x' = \frac{y'^2}{4a}$ ;

hence, by substitution in (1), we obtain  $\frac{4a}{y} = \frac{y'}{a}$ .

Again,  $\frac{SQ}{AS} = \frac{PM}{AM}$ ;

or,  $\frac{SQ}{a} = \frac{y}{x} = \frac{4a}{y} = \frac{y'}{a} = \frac{P'M'}{a}$ ;  $\therefore SQ = P'M'$ ;

but the triangles  $QSE$  and  $PM'E$  are similar;  $\therefore QE = EP'$ .

8. Let  $AM = x$ ,  $PM = y$ ; draw  $QN'$  perpendicular to  $DS$ , and let  $SN' = h$ ,  $QN' = k$ . By similar triangles,

$$\frac{SN'}{QN'} = \frac{SM}{PM}, \text{ or } \frac{h}{k} = \frac{a-x}{y},$$

$$\frac{DS}{SN} = \frac{DM}{PM}, \text{ or } \frac{2a}{k} = \frac{a+x}{y};$$

$$\therefore \frac{2a+h}{k} = \frac{2a}{y}, \text{ and } \frac{2a-h}{k} = \frac{2x}{y};$$

hence, by multiplication,  $\frac{4a^2 - h^2}{k^2} = \frac{4ax}{y^2} = 1$ ,

$$\text{or, } h^2 + k^2 = 4a^2,$$

which is the equation to a circle whose centre is S, and radius  $= 2a = DS$ .

9. Let  $y^2 = 4ax$  be the equation to the parabola. The equation to the tangent, in terms of the tangent of the angle which the line makes with the axis, is

$$y' = mx' + \frac{a}{m}, \text{ or } m^2 - \frac{y'}{x'} m + \frac{a}{x'} = 0;$$

solving this quadratic, we obtain

$$2mx' = y' \pm \sqrt{y'^2 - 4ax'};$$

$$\text{but } x' = \frac{y'}{m} - \frac{a}{m^2}, \text{ or } 2mx' = 2y' - \frac{2a}{m};$$

$$\therefore 2y' - \frac{2a}{m} = y' \pm \sqrt{y'^2 - 4ax'};$$

$$\text{hence } \frac{1}{m} = \frac{y'}{2a} \mp \frac{1}{2a} \sqrt{y'^2 - 4ax'}.$$

The difference of these two cotangents is

$$\frac{1}{a} \sqrt{y'^2 - 4ax'} = d, \text{ a constant};$$

$$\therefore y^2 - 4ax = a^2 d^2, \text{ or } y^2 = 4a(x + \frac{1}{4}ad^2),$$

which is the eqn. to a parabola whose latus rectum is  $4a$ , and vertex at a distance  $= \frac{1}{4}ad^2$  from the origin.

### EXERCISES [G].

$$1. \frac{\sin^2 \theta}{\cos^2 \theta} = \frac{9}{1}; \therefore \frac{\sin^2 \theta}{\sin^2 \theta + \cos^2 \theta} = \frac{9}{10};$$

$$\therefore \sin \theta = \frac{3}{\sqrt{10}}, \cos \theta = \sqrt{1 - \frac{9}{10}} = \frac{1}{\sqrt{10}} \sqrt{10}.$$

For  $x$  put  $x' \cos \theta - y' \sin \theta$ , and for  $y$  put  $x' \sin \theta + y' \cos \theta$   
then  $3xy - 4x^2 = a^2$  will become  $\frac{1}{2}x'^2 - \frac{9}{2}y'^2 = a^2$ ,

$$\text{or, } x'^2 - 9y'^2 = 2a^2.$$

2. Let  $x' = x \cos \theta - y \sin \theta$ , and  $y' = x \sin \theta + y \cos \theta$ ,  
 or  $x' = \frac{1}{2}\sqrt{2}(x-y)$ , and  $y' = \frac{1}{2}\sqrt{2}(x+y)$ ;  
 $\therefore x'y' = \frac{1}{2}(x^2 - y^2) = \frac{1}{2}a^2$ .

3. Here  $A=4$ ,  $B=12$ ,  $C=9$ ;  $\therefore \tan 2\theta = -\frac{1}{2}$ ,  
 $\therefore \tan \theta = \frac{3}{2}$ ,  $\sin \theta = \frac{3}{\sqrt{13}}$ ,  $\cos \theta = \frac{2}{\sqrt{13}}$ ;

$$\text{hence } x'^2 = \frac{13}{100}, \text{ or } x' = \pm \frac{1}{10}\sqrt{13};$$

the equation therefore represents two parallel straight lines, each inclined to the axis of  $x$  at an angle whose tangent is  $-\frac{3}{2}$ , and whose distance from each other is  $\pm \frac{1}{10}\sqrt{13}$ .

4. Here  $A=1$ ,  $B=1$ ,  $C=1$ ;  $\therefore \tan 2\theta = 1+0=\infty$ , the tang. of  $90^\circ$ ;  $\therefore \theta = 45^\circ$ ,  $\sin \theta = \cos \theta = \frac{1}{2}\sqrt{2}$ ;

$$\text{hence } 3x'^2 + y'^2 = 2, \text{ or } \frac{x'^2}{\frac{2}{3}} + \frac{y'^2}{2} = 1,$$

the equation to an ellipse whose axes are  $2a = \frac{2}{3}\sqrt{6}$ , and  $2b = 2\sqrt{2}$ .

5. Here we have  $A=9$ ,  $B=-30$ ,  $C=25$ ,  $D=21$ ,  $E=-35$ ,  $F=10$ ; and since  $B^2 - 4AC = 0$ , we cannot get rid of the simple powers of  $x$  and  $y$ ; but we have  $\tan 2\theta = \frac{1}{2}$ ;

$$\therefore \tan \theta = \frac{3}{4}, \sin \theta = \frac{3}{5}, \cos \theta = \frac{4}{5}; \text{ and applying these}$$

in the usual way, the given equation is transformed to  $34y'^2 - 7\sqrt{34}y' + 10 = 0$ , an equation yielding two numerical values of  $y'$ . So that the given equation represents two parallel straight lines. It will be found to be the product of  $5y - 3x - 2 = 0$  and  $5y - 3x - 5 = 0$ , each of which represents a straight line the tangent of whose inclination to the axis of  $x$  is  $\frac{3}{5}$ .

6. Here we have  $A=1$ ,  $B=-6$ ,  $C=1$ ,  $D=-6$ ,  $E=2$ ; and since  $B \div (A-C)$  is  $=\infty$ , we can get rid of the simple powers of  $x$  and  $y$ ; accordingly, we obtain  $h=0$ ,  $k=-1$ ; and putting  $y=y'-1$ , the given equation is transformed to

$x'^2 - 6x'y' + y'^2 + 4 = 0$ . Now,  $\tan 2\theta$  being  $=\infty$ , we have  $\theta = 45^\circ$ ,  $\sin \theta = \cos \theta = \frac{1}{\sqrt{2}}$ ; and transforming we obtain

$$x''^2 - 2y''^2 = 2, \text{ or } \frac{x''^2}{2} - \frac{y''^2}{1} = 1,$$

the equation to an hyperbola, whose axes are  $2a = 2\sqrt{2}$  and  $2b = 2$ .

7. Here we have  $x^2 + 2xy - y^2 - 2 = 0$ , where  $A = 1$ ,  $B = 2$ ,  $C = -1$ ,  $F = -2$ ;  $\tan 2\theta = 1$ ;  $\therefore \tan \theta = \sqrt{2} - 1$ ,  $\sin \theta = \frac{1}{\sqrt{2}}\sqrt{2 - \sqrt{2}}$ ,  $\cos \theta = \frac{1}{\sqrt{2}}\sqrt{2 + \sqrt{2}}$ . Accordingly, the equation will assume the form  $x'^2 - y'^2 = \sqrt{2}$ , the eqn. to a rectangular hyperbola, where  $a^2 = \sqrt{2}$ , or  $a = b = 2^{\frac{1}{4}}$ .

8. Here  $B^2 - 4AC = 0$ .  $\tan 2\theta = -2 + 0 = \infty$ ;  $\therefore \theta = 45^\circ$ ,  $\sin \theta = \cos \theta = \frac{1}{\sqrt{2}}$ . Accordingly, the eqn. will assume the form

$$y'^2 - 2\sqrt{2}x' + 2\sqrt{2}y' + 8 = 0,$$

where  $C' = 1$ ,  $D' = -2\sqrt{2}$ ,  $E' = 2\sqrt{2}$ ,  $F = 8$ .

$$\left(y' + \frac{E'}{2C'}\right)^2 = -\frac{D'}{C'}\left(x' + \frac{F}{D'} - \frac{E'^2}{4C'D'}\right),$$

$$\text{or } (y' + \sqrt{2}) = 2\sqrt{2}(x' - 1\frac{1}{2}\sqrt{2});$$

$$\therefore y'' = 2\sqrt{2}x'',$$

the equation to a parabola; the coordinates of the vertex being  $1\frac{1}{2}\sqrt{2}$  and  $-\sqrt{2}$ .

9. Here we have  $x^2 - 2xy + y^2 - 2x - 2y - 3 = 0$ ;

$$B^2 - 4AC = 0. \quad \tan 2\theta = \infty, \quad \therefore \theta = 45^\circ,$$

$\sin \theta = \cos \theta = \frac{1}{\sqrt{2}}$ . Accordingly, the equation is transformed to  $2y'^2 - 2\sqrt{2}x' - 3 = 0$ , where  $C' = 2$ ,  $D' = -2\sqrt{2}$ ,  $F = -3$ ; we hence obtain

$$y'^2 = \sqrt{2}\left(x' + \frac{3}{2\sqrt{2}}\right), = \sqrt{2}\left(x' + \frac{3}{4}\sqrt{2}\right);$$

the eqn. to a parabola; the coordinates of the vertex being  $-\frac{3}{4}\sqrt{2}$  and 0; latus rectum  $= \sqrt{2}$ .

10. Here  $h = \frac{1}{2}$ ,  $k = -\frac{1}{2}$ ; transformed equation,  $3x'^2 - 2x'y' + y'^2 - 4\frac{1}{2} = 0$ ;  $\tan 2\theta = -1$ ,  $\therefore 2\theta = 135^\circ$ ,  $\theta = 67\frac{1}{2}^\circ$ ;  $\sin \theta = \frac{1}{\sqrt{2}}\sqrt{2 + \sqrt{2}}$ ,  $\cos \theta = \frac{1}{\sqrt{2}}\sqrt{2 - \sqrt{2}}$ ; whence, by further transformation,

$$(4 - 2\sqrt{2})x'^2 + (4 + 2\sqrt{2})y'^2 = 9,$$

the eqn. to an ellipse; the coordinates of the centre  $\frac{1}{2}$  and  $-\frac{1}{2}$ , and the axes,  $a=1\frac{1}{2}\sqrt{2+\sqrt{2}}$ ,  $b=1\frac{1}{2}\sqrt{2-\sqrt{2}}$ .

11. Take C, the middle point of BA as origin; BA = 2n, CM = x, PM = y;

$$\tan A = \frac{y}{n-x}, \tan B = \frac{y}{n+x};$$

$$\tan(A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B} = \frac{2xy}{n^2 - x^2 + y^2} = \tan 30^\circ = \frac{1}{3}\sqrt{3};$$

$$\text{whence } x^2 + 2\sqrt{3}xy - y^2 - n^2 = 0.$$

$$\tan 2\theta = \sqrt{3} = \tan 60^\circ, \therefore \theta = 30^\circ, \sin \theta = \frac{1}{2}, \cos \theta = \frac{1}{2}\sqrt{3};$$

hence the transformed eqn. is  $x'^2 - y'^2 = \frac{1}{2}n^2$ ;

or the locus of P is an equilateral hyperbola, whose axes are each  $=\frac{1}{2}\sqrt{2}n$ , and whose transverse axis makes with the base of the triangle an angle  $=30^\circ$ .

12. Let ACB be one of the equal triangles on the base AB. Draw AD, BE perpendiculars to BC, AC; and through C draw the straight line CPM to meet AB.

Let A be the origin of coordinates, AM = x', CM = y'; and put AB = c.

The eqn. to AC, passing through the points (0, 0) and (x', y'), is

$$y = \frac{y'}{x'}x.$$

The eqn. to BC, passing through the points (c, 0) and (x', y'), is



$$y = \frac{y'}{x' - c}(x - c).$$

Now, BE being perpend. to AC, and passing through the point (c, 0), its eqn. is

$$y = -\frac{x'}{y'}(x - c).$$

The eqn. to AD, a perpend. on BC from the origin, is

$$y = -\frac{x' - c}{y'}x.$$



Now, at the point P where these perpendiculars intersect, the ordinate is common ;

$$\therefore \frac{x'}{y'}(x-c) = \frac{x'-c}{y'}x ;$$

whence  $x=x'$ , that is,  $x$ , the abscissa of the point P, is also the abscissa of the point C ;  $\therefore$  CPM is perpend. to AB.

Now, to find the locus of P for all triangles having the same base and altitude,

let  $AM=x$ ,  $PM=y$ ,  $CM=p$ , the constant altitude.

$$\cot ABD = \tan DAB = \frac{PM}{AM} = \frac{y}{x} ;$$

$$\tan ABD = \frac{CM}{BM} = \frac{p}{c-x} ;$$

by mult<sup>n</sup>,  $1 = \frac{py}{cx-x^2}$ , or  $x^2 + py - cx = 0$  ;

where  $C'=1$ ,  $D'=p$ ,  $E'=-c$ .

$$\left(x' - \frac{c}{2}\right)^2 = -p\left(y' - \frac{c^2}{4p}\right),$$

which is the equation to a parabola having its axis parallel to that of  $y$ , the coordinates of its vertex being  $\frac{c}{2}$  and  $\frac{c^2}{4p}$ , and its latus rectum  $=p$ .

LONDON

PRINTED BY SPOTTISWOODE AND CO.

NEW-STREET SQUARE

## *Works by the same Author.*

---

**MODERN ARITHMETIC.** A Treatise adapted for School Work and for Private Study. Containing numerous improvements in aid of the Preparation of Candidates for Public Examinations....12mo. [*Nearly ready.*]

**EXERCISES in the FIRST FOUR RULES of ARITHMETIC,** constructed for the Application of New Artificial Tests, by which the Teacher may expeditiously ascertain the Correctness of the Results. Third Edition.....12mo. 6d.

**SOLUTIONS of QUESTIONS in ARITHMETIC and BOOK-KEEPING** used in the Civil Service Examinations of 1862, and published in the Appendix to the Eighth Report of the Commissioners: with a SUPPLEMENT containing Examples in Account-States, &c. ....12mo. 1s. 6d.

**MERCANTILE EXERCISES in ADDITION of MONEY and in the CALCULATION of PER-CENTAGES;** including numerous Examples of a General and Simple Method of treating Per-Centage Problems, with Answers to the Exercises .....12mo. 1s. 3d.

**PROGRESSIVE EXERCISES in BOOK-KEEPING by DOUBLE ENTRY,** including Account-States, Partnership Accounts, Private Journal, and Ledger, &c. ....12mo. 1s. 6d.

**EXAMINATION-QUESTIONS in BOOK-KEEPING by DOUBLE ENTRY,** preceded by full Directions for the True Stating of Dr. and Cr.; with ANSWERS, providing the means of solving all the Problems and Exercises relating to Account-Books in the Civil Service and other Examination-Papers .....12mo. 2s. 6d.

**HUNTER'S EXAMINATION-QUESTIONS and DIRECTIONS,** as above, separated from the ANSWERS.....12mo. 1s.

**RULED PAPER** for the various Forms of Account-Books required in HUNTER'S Examination-Questions in Book-keeping, 5 sorts, price 1s. 6d. per Quire.

**EXAMINATION-QUESTIONS on COLEN'S *ELEMENTS of ALGEBRA*, PART I.** ....12mo. 2s. 6d.

**TREATISE on LOGARITHMS:** With Copious Tables of Selected Logarithms; explaining simply the Nature and use of Logarithms and Logarithmic Tables, the Principles and Methods of their Construction, and their Application: With numerous Examples and Exercises .....16mo. 1s. KEY, 9d.

**ELEMENTS of MENSURATION,** simplified for the use of Beginners; with numerous Original Problems and Progressive Exercises. New Edition, Revised and Stereotyped .....18mo. 9d.—KEY, 9d.

**ELEMENTS of PLANE TRIGONOMETRY,** for Beginners; with numerous Problems, and Tables of all the Natural Sines, &c. required for the Solution of the Exercises .....18mo. 1s.—KEY, 9d.



**TEXT-BOOK of ENGLISH GRAMMAR:** a Treatise on the Etymology and Syntax of the English Language; including Exercises in Parsing and Punctuation: an Etymological Vocabulary of Grammatical Terms; and a copious List of the principal Works on English Grammar. New Edition, and thoroughly Revised .....12mo. 2s. 6d.

**EXERCISES in ENGLISH PARSING,** Progressively arranged and adapted to the Author's *Text-Book of English Grammar*; with Questions suggesting a Course of Oral Instruction for Junior Pupils. Tenth Edition .....12mo. 6d.

**PARAPHRASING and ANALYSIS of SENTENCES,** simplified for the use of Schools, forming a Manual of Instruction and Exercise for the use of Students, Teachers, &c. ....12mo. 1s. 3d.—KEY, 1s. 3d.

**SCHOOL MANUAL of LETTER WRITING:** Containing numerous Models of Letters on Commercial and other subjects; with Exercises in Epistolary Composition, Rules of Punctuation, Explanations of Abbreviated Titles, Commercial Terms, &c. Second Edition .....12mo. 1s. 6d.

**INTRODUCTION to the WRITING of PRECIS or DIGESTS,** as applicable to Narratives of Facts or Historical Events, Correspondence, Evidence, Official Documents, and General Composition: with numerous Examples and Exercises. Fourth Edition.....12mo. 2s.—KEY, 1s.

**JOHNSON'S *RASSELAS*,** with Introductory Remarks, Explanatory and Grammatical Annotations, Specimens of Interrogative Lessons, Answers to Examination Questions, &c. ....12mo. 2s. 6d.

**BACON'S *ADVANCEMENT of LEARNING*,** Annotated Summary of Books I. and II. with numerous Extracts from the Work and Specimens of Examination-Questions.....12mo. 2s.

**EXAMINATION-QUESTIONS on the FIRST TWO BOOKS of MILTON'S *PARADISE LOST*,** and on SHAKSPEARE'S *MERCHANT OF VENICE*, for the use of Middle-Class Candidates .....12mo. 1s.

**SELECT PLAYS of SHAKSPEARE,** with Explanatory and Illustrative Notes, Critical Remarks, and other Aids to a thorough understanding of each Drama. Edited for the use of Schools and Pupils preparing for Examination .....12mo. 2s. 6d. each Play.

HENRY VIII.  
JULIUS CÆSAR.

MERCHANT of VENICE.  
HAMLET.

THE TEMPEST.  
KING LEAR.

**BOOKS I. and II. of MILTON'S *PARADISE LOST*:** with a Prose Translation or Paraphrase, the Parsing of the more Difficult Words, Specimens of Analysis, and numerous Illustrative Notes, for the use of Candidates.....12mo. price 1s. 6d. each Book.

**MILTON'S *COMUS*, *L'ALLEGRO*, and *IL PENSEROSO*,** with numerous Notes, critical, illustrative, and explanatory: adapted for use in Training Colleges and Schools .....12mo. 1s. 6d.






# GLEIG'S SCHOOL SERIES.

New Series of Elementary School-Books; each Book (in most instances) complete in itself, price NINEPENCE. Intended to comprise a complete Course of Elementary Education. Projected and Edited by the Rev. G. R. GLEIG, M.A. Chaplain-General to Her Majesty's Forces. Assisted by

WALTER M'LEOD, F.R.G.S. | THE REV. J. HUNTER, M.A. | THOMAS TATE, F.R.A.S.  
 WILLIAM HUGHES, F.R.G.S. | DR. R. J. MANN, M.R.C.S.E. | A. K. ISBISTER, M.A.  
 MR. JAMES OWEN. | PROF. R. W. BROWNE M.A. | W. J. REYNOLDS, M.A. ETC.



	s.	d.
FIRST SCHOOL-BOOK to TEACH READING and WRITING.....	0	6
SECOND SCHOOL-BOOK to TEACH READING and SPELLING ....	0	9
A GRADUATED SERIES of NINE COPY-BOOKS .....	0	3
SIMPLE TRUTHS from SCRIPTURE .....	0	6
EXPLANATORY ENGLISH GRAMMAR, 9d.....	0	1
HISTORY of the ENGLISH LANGUAGE.....	0	9
The CHILD'S FIRST GEOGRAPHY .....	0	9
GEOGRAPHY of the BRITISH EMPIRE .....	0	9
GENERAL GEOGRAPHY, 9d.....	0	9
QUESTIONS ON GENERAL GEOGRAPHY .....	0	9
HAND-ATLAS of GENERAL GEOGRAPHY, 29 full-coloured Maps, sewed, 2s. 6d. ....	3	0
half-bound .....	3	0
CLASS-ATLAS of PHYSICAL GEOGRAPHY, 30 full-coloured Maps, Sections, and Diagrams, sewed, 2s. 6d. ....	3	0
half-bound .....	3	0
BOWMAN'S <i>Questions on M'Leod's Physical Atlas</i> .....	1	0
PHYSICAL ATLAS of GREAT BRITAIN and IRELAND, fcp. 4to.....	7	6
SACRED HISTORY, 2s. ; or in Two Parts .....	0	9
each .....	0	9
each .....	0	9
HISTORY of the BRITISH COLONIES .....	0	9
HISTORY of BRITISH INDIA .....	0	9
HISTORICAL QUESTIONS, PART I. on the above Four Histories .....	0	9
HISTORY of FRANCE .....	0	9
HISTORY of GREECE, 9d.....	0	9
HISTORY of ROME, .....	0	9
BOOK of BIOGRAPHY .....	0	9
ASTRONOMY and the USE of the GLOBES .....	0	9
NAVIGATION and GREAT CIRCLE SAILING.....	1	0
ELEMENTS of EUCLID, 9d.....	1	0
PRACTICAL GEOMETRY, .....	1	0
ORTHOGRAPHIC PROJECTION and ISOMETRICAL DRAWING ....	1	0
ELEMENTS of MENSURATION, 9d.....	0	9
KEY, .....	0	9
ELEMENTS of PLANE TRIGONOMETRY, 1s.....	0	9
KEY, .....	0	9
TREATISE on LOGARITHMS, with TABLES, 1s. ....	0	9
KEY, .....	0	9
A MANUAL of ARITHMETIC .....	9	
BOOK-KEEPING by SINGLE and DOUBLE ENTRY .....	0	9
A Set of Eight Account-Books, adapted to the above, price 6d. each.		
ELEMENTS of ALGEBRA, 9d. ANSWERS to PROBLEMS in ditto, 3d.		
Full KEY .....	1	0
HYDROSTATICS, HYDRAULICS, and PNEUMATICS .....	0	9
The BOOK of HEALTH.....	0	9
BOOK of DOMESTIC ECONOMY .....	0	9
ELECTRICITY, 9d.....	0	9
LIGHT and HEAT, .....	0	9
MAGNETISM, VOLTAIC ELECTRICITY, and ELECTRO-DYNAMICS, .....	0	9
EXPERIMENTAL CHEMISTRY .....	9	
MECHANICS and the STEAM-ENGINE.....	0	9
NATURAL HISTORY for BEGINNERS, 2s. ; or in 2 Parts.....	0	9
each .....	0	9

London: LONGMANS, GREEN, and CO. Paternoster Row.

